


11p

**NASA CONTRACTOR
REPORT**



NASA CR-50

NASA CR-50

N64-21253

Code 1

cat-30

220

ROCKET BOOSTER CONTROL

**A SUFFICIENT CONDITION IN
OPTIMAL CONTROL**

by E. B. Lee

Prepared under Contract No. NASw-563 by
MINNEAPOLIS-HONEYWELL REGULATOR COMPANY
Minneapolis, Minnesota

for

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION • WASHINGTON, D. C. • MAY 1964

ROCKET BOOSTER CONTROL

A SUFFICIENT CONDITION IN OPTIMAL CONTROL

By E. B. Lee

**Prepared under Contract No. NASw-563 by
MINNEAPOLIS-HONEYWELL REGULATOR COMPANY
Minneapolis, Minnesota**

**This report is reproduced photographically
from copy supplied by the contractor.**

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

**For sale by the Office of Technical Services, Department of Commerce
Washington, D. C. 20230 -- Price \$0.50**

TABLE OF CONTENTS

ABSTRACT	1
INTRODUCTION	1
DEVELOPMENT	3
REMARKS	7
CONCLUSIONS	7
REFERENCES	8

A SUFFICIENT CONDITION IN OPTIMAL CONTROL*

By E. B. Lee[†]

ABSTRACT

21253

A theorem is proven which covers most of the known cases where L. S. Pontriagins' Maximum Principle is a sufficient, as well as a necessary, condition for optimal control.

Author

INTRODUCTION

Consider the system

$$\dot{x}^0 = f^0(x, t) + h^0(u, t) \quad (1)$$

$$\dot{x} = A(t)x + h(u, t) \quad (2)$$

with $x(t_0) = x_0$ and $x^0(t_0) = 0$. Here f^0 , h^0 , A , and h are continuous in all arguments. x is the system state, an n vector; u is the control, an m vector. x^0 is a scalar variable which measures the quality of control.

If $u(s)$ is any control function on the interval $[t_0, t]$ we will write the corresponding response of equations (1) and (2) as $\hat{x}_u(t) = (x_u^0(t), x_u(t))$. The control u is restricted to a set $\Omega \subset R^m$. It is assumed that either Ω is compact or that h , and h^0 are such that

$$\max_{u \in \Omega} \{ \lambda \cdot h(u, t) + \lambda^0 h^0(u, t) \}$$

exists for each $t \in [t_0, T]$ and $\hat{\lambda} = (\lambda^0, \lambda) \in R^{n+1}$ with $\lambda^0 < 0$.

It is further assumed that $f^0(x, t)$ is a single-valued, convex function of x for each $t \in [t_0, T]$, that is,

* Prepared under contract NASw-563 for the NASA

† Research Consultant, Minneapolis-Honeywell Regulator Company, Minneapolis, Minnesota.

$$\frac{\partial f^0}{\partial x}(x, t) \cdot (\omega - x) + f^0(x, t) \leq f^0(\omega, t)$$

for all x and $\omega \in \mathbb{R}^n$ and $t \in [t_0, T]$.

The following definitions are needed:

DEFINITION 1. $u^*(s)$ on $[t_0, T]$ is an extremal control if there exists $\lambda(t_0) = (\lambda_0^1, \lambda_0^2, \dots, \lambda_0^n)$ and $\lambda^0 = \text{constant} < 0$, such that

$$\lambda^0 h^0(u^*(s), s) + \lambda(s) \cdot h(u^*(s), s) = \max_{u \in \Omega} \{ \lambda^0 h^0(u, s) + \lambda(s) \cdot h(u, s) \}$$

with

$$\dot{\lambda} = -A'(t)\lambda - \lambda^0 \frac{\partial f^0}{\partial x}(x_{u^*}(t), t), \quad (' \text{ denotes transpose}).$$

Here $x_{u^*}(t)$ is the response corresponding to $u^*(s)$, $t_0 \leq s \leq t \leq T$.

DEFINITION 2. The control $u(s)$ is allowable if it is a measurable real valued vector function with range in Ω on $[t_0, T]$.

DEFINITION 3. The set of attainability $K(T, x_0)$ is the collection of end points of the responses $\hat{x}_u(t) = (x_u^0(t), x_u(t))$ which initiate at $(0, x_0)$ for all allowable controls $u(s)$ on $[t_0, T]$.

The problem of optimal control studied here is to select allowable controls $u(s)$ which "steer" the response $x_u(t)$ from the initial point x_0 at time t_0 to a prescribed target set G at time $T < \infty$ and minimize the cost functional of control $C(u) = g(x(T)) + x^0(T)$. Here g is a continuously differentiable function of x . An allowable control which provides an absolute minimum for $C(u)$ amongst the set of all allowable controls which steer $x_u(t)$ from x_0 to G is called an optimal control. Note, the free end point problem results when $G = \mathbb{R}^n$.

DEVELOPMENT

A basic inequality is established and then the main theorem is proven.

LEMMA. Let $u^*(s)$, $t_0 \leq s \leq T$, be an allowable extremal control with corresponding response $\hat{x}_{u^*}(t)$ which initiates at $\hat{x}(t_0) = (0, x(t_0))$, then $\hat{\lambda}(T) \cdot \hat{x}_{u^*}(T) \geq \hat{\lambda}(T) \cdot \hat{\omega}$ for $\lambda^0 < 0$ and all $\hat{\omega} \in K(T, x_0)$.

PROOF. Consider

$$\begin{aligned} \frac{d(\hat{\lambda} \cdot \hat{x}_{u^*})}{dt} &= \lambda^0 \dot{x}_{u^*}^0 + \dot{\lambda} \cdot x_{u^*} + \lambda \cdot \dot{x}_{u^*} \\ &= \lambda^0 (f^0(x_{u^*}, t) + h^0(u^*, t)) + \\ &\quad (-A'(t)\lambda - \lambda^0 \frac{\partial f^0}{\partial x}(x_{u^*}, t)) \cdot x_{u^*} \\ &\quad + \lambda \cdot (A(t)x_{u^*} + h(u^*, t)). \end{aligned}$$

Upon integrating both sides between t_0 and T we obtain

$$\begin{aligned} \lambda^0 x_{u^*}^0(T) + \lambda(T) \cdot x_{u^*}(T) - \lambda(t_0) \cdot x(t_0) = \\ \int_{t_0}^T \left\{ \lambda^0 (f^0(x_{u^*}(t), t) - \frac{\partial f^0}{\partial x}(x_{u^*}(t), t) \cdot x_{u^*}(t)) \right. \\ \left. + \lambda^0 h^0(u^*(t), t) + \lambda(t) \cdot h(u^*(t), t) \right\} dt \end{aligned}$$

Let $x_u(t)$ be any other response with initial value $x_0 = x(t_0)$ for which we calculate

$$\begin{aligned} \lambda^0 x_u^0(T) + \lambda(T) \cdot x_u(T) - \lambda(t_0) \cdot x(t_0) = \\ \int_{t_0}^T \left\{ \lambda^0 (f^0(x_u(t), t) - \frac{\partial f^0}{\partial x}(x_{u^*}(t), t) \cdot x_u(t)) \right. \\ \left. + \lambda^0 h^0(u(t), t) + \lambda(t) \cdot h(u(t), t) \right\} dt. \end{aligned}$$

But $\lambda^0 h^0(u^*(t), t) + \lambda(t) \cdot h(u^*(t), t) \geq \lambda^0 h^0(u(t), t) + \lambda(t) \cdot h(u(t), t)$.

$$\begin{aligned} \text{Thus if } \lambda^0(f^0(x_{u*}(t), t) - \frac{\partial f^0}{\partial x}(x_{u*}(t), t) \cdot x_{u*}(t)) \geq \lambda^0(f^0(x_u(t), t) \\ - \frac{\partial f^0}{\partial x}(x_{u*}(t), t) \cdot x_u(t)) \end{aligned}$$

we obtain the desired inequality. This is certainly true if $\lambda^0 < 0$ and

$$\begin{aligned} \frac{\partial f^0}{\partial x}(x_{u*}(t), t) \cdot x_{u*}(t) - f^0(x_{u*}(t), t) \geq \frac{\partial f^0}{\partial x}(x_{u*}(t), t) \cdot x_u(t) - \\ - f^0(x_u(t), t), \end{aligned}$$

which is the convexity condition on f^0 .

Thus we have

$$\lambda^0 x_{u*}^0(T) + \lambda(T) \cdot x_{u*}(T) \geq \lambda^0 x_u^0(T) + \lambda(T) \cdot x_u(T) \text{ or}$$

$$\hat{\lambda}(T) \cdot \hat{x}_{u*}(T) \geq \hat{\lambda}(T) \cdot \hat{\omega} \text{ all } \hat{\omega} \in K(T, x_0) \text{ and the lemma}$$

is established.

The basic inequality of the lemma enables us to establish the sufficiency of the maximum principle in a number of cases. These results are summarized as a theorem:

THEOREM.

A) Consider the cost functional of control $C(u) = x^0(T)$ and as target set G a point x_1 . Let $u^*(s)$, $t_0 \leq s \leq T$, be an allowable extremal control which steers the corresponding response $x_{u*}(t)$ from x_0 at t_0 to x_1 at T , then $u^*(s)$ is an optimal control.

B) Consider the cost functional $C(u) = g(x(T)) + x^0(T)$ with $g(x)$ a convex function of x and consider the target set $G = R^n$, (this is the free end point problem). Let $x(t_0) = x_0$. Then $u^*(s)$, $t_0 \leq s \leq T$, is an optimal control if it is an allowable extremal control with $\hat{\lambda}(T) = (-1, -\frac{\partial g}{\partial x}(x_{u*}(T)))$, (The condition on $\hat{\lambda}(T)$ is called a transversality condition).

C) Consider the cost functional $C(u) = x^0(T)$ and the convex, closed, target set $G = \{x | \gamma(x) \leq c\} \subset \mathbb{R}^n$, where γ is differentiable and c a constant. Let $u^*(s)$, $t_0 \leq s \leq T$, be an allowable extremal control which steers $x_{u^*}(t)$ from x_0 at t_0 to $x_1 \in G$ at T with $\lambda(T)$ an interior normal[†] to G at $x_{u^*}(T)$ on ∂G , then $u^*(s)$ is optimal if such a control exists. (If there is no such $u^*(s)$ then the minimum may occur interior to G in which case B) applies with $\hat{\lambda}(T) = (-1, 0, 0 \dots 0)$ and if G is just one point part A) is obtained.)

PROOF:

A) From the lemma

$$\hat{\lambda}(T) \cdot \hat{x}_{u^*}(T) \geq \hat{\lambda}(T) \cdot \hat{\omega} \text{ for } \lambda^0 < 0 \text{ all } \hat{\omega} \in K(T, x_0).$$

Thus

$$\lambda(T) \cdot x_{u^*}(T) + \lambda^0 x_{u^*}^0(T) \geq \lambda(T) \cdot x_u(T) + \lambda^0 x_u^0(T).$$

But, comparing only those responses that end at x_1 , that is, those for which $x_{u^*}(T) = x_u(T) = x_1$, the basic inequality becomes

$$\lambda^0 x_{u^*}^0(T) \geq \lambda^0 x_u^0(T).$$

Since $\lambda^0 < 0$ we have $C(u^*) = x_{u^*}^0(T) \leq x_u^0(T) = C(u)$ and therefore $u^*(s)$ is optimal.

B) With $\hat{\lambda}(T) = (-1, \frac{-\partial g}{\partial x}(x_{u^*}(T)))$ the inequality of the lemma is

$$\frac{-\partial g}{\partial x}(x_{u^*}(T)) \cdot x_{u^*}(T) - x_{u^*}^0(T) \geq \frac{-\partial g}{\partial x}(x_{u^*}(T)) \cdot x_u(T) - x_u^0(T)$$

[†] λ is an interior normal to G at x on ∂G if λ is orthogonal to a support plane of G at x and is directed into the halfspace containing G . Thus G need not have an interior to have interior normals. Note if G does not have an interior we can still approximate it by a $\gamma(x)$.

Adding and subtracting $g(x_{u*}(T))$ on the left side and $g(x_u(T))$ on the right side the last inequality becomes

$$\begin{aligned} -x_{u*}^0(T) - g(x_{u*}(T)) + g(x_{u*}(T)) - \frac{\partial g}{\partial x}(x_{u*}(T)) \cdot x_{u*}(t) &\geq \\ -x_u^0(T) - g(x_u(T)) + g(x_u(T)) - \frac{\partial g}{\partial x}(x_{u*}(T)) \cdot x_u(T). \end{aligned}$$

But,

$$\frac{\partial g}{\partial x}(x_{u*}(T)) \cdot (x_{u*}(T) - x_u(T)) + g(x_u(T)) \geq g(x_{u*}(T))$$

if g is a convex function of x .

Therefore

$$\begin{aligned} -C(u^*) &= -x_{u*}^0(T) - g(x_{u*}(T)) \geq \\ -x_u^0(T) - g(x_u(T)) &= -C(u), \end{aligned}$$

or $C(u^*) \leq C(u)$. Hence part B) is established.

C) Assume for simplicity that γ was picked to be a convex function on G with $\partial G = \{x | \gamma(x) = c\}$. Consider only boundary points $x_{u*}(T)$ where it is required that $\lambda(T) = k \left\{ -\frac{\partial \gamma}{\partial x}(x_{u*}(T)) \right\}$ in order for $\lambda(T)$ to be an interior normal to G at $x_{u*}(T)$ on ∂G , (let $k = 1$).

The inequality of the lemma can then be written

$$\lambda(T) \cdot x_{u*}(T) = \lambda^0 x_{u*}^0(T) + \left\{ -\frac{\partial \gamma}{\partial x}(x_{u*}(T)) \right\} \cdot x_{u*}(T) \geq$$

$$\lambda(T) \cdot x_u(T) = \lambda^0 x_u^0(T) + \left\{ -\frac{\partial \gamma}{\partial x}(x_{u*}(T)) \right\} \cdot x_u(T)$$

If $x_{u*}(T)$ is on the boundary of G it is also true that

$\gamma(x_{u*}(T)) = c \geq \gamma(x_u(T))$ for all allowable responses $x_u(T) \in G$.

Adding the last two inequalities we obtain

$$\begin{aligned} \lambda^0 x_{u*}^0(T) - \frac{\partial \gamma}{\partial x}(x_{u*}(T)) \cdot x_{u*}(T) + \gamma(x_{u*}(T)) &\geq \\ \lambda^0 x_u^0(T) - \frac{\partial \gamma}{\partial x}(x_{u*}(T)) \cdot x_u(T) + \gamma(x_u(T)). \end{aligned}$$

But again if $x_{u*}(T)$ is on ∂G and $x_u(T)$ is in the convex set G we

have

$$\frac{\partial \gamma}{\partial x}(x_{u^*}(T)) \cdot [x_{u^*}(T) - x_u(T)] + \gamma(x_u(T)) \geq \gamma(x_{u^*}(T))$$

and therefore

$$C(u^*) \leq C(u). \quad \text{Q.E.D.}$$

REMARKS

If the set of attainability is closed there will exist optimum control provided there is at least one control that steers the response to the desired end point $x_1 \in G$, assuming G is also closed. The property of closure is discussed in reference 2 in which a bibliography and discussion of cases are presented. The set of attainability is also known to be closed if

$h(u, t) = B(t)u$, $f^0(x, t) = x \cdot W(t)x$ and $h^0(u, t) = u \cdot U(t)u$, for $W(t)$, $U(t)$ positive definite on $[t_0, T]$.

When the set of attainability is closed, in the above case, the inequality of the lemma establishes that its lower (exterior normal with $\lambda^0 < 0$) surface is convex. For if it was otherwise we would be led to a contradiction of the maximum principle. Note that the transversality condition (reference 1) follows from the established inequality of the lemma since $\hat{\lambda}(T)$ must be an exterior normal of $\bar{K}(T, x_0)$ at the corresponding response end point, $\hat{x}_{u^*}(T)$.

CONCLUSIONS

It has been proven the Maximum Principle is often a sufficient as well as necessary condition for optimal control.

REFERENCES

1. Pontriagin, L. S., Boltyanskii, V. G., Gamkrelidze, R. V., and Mischenko, E. R., "The Mathematical Theory of Optimal Control Processes", Interscience, 1962
2. Neustadt, L. W., "The Existence of Optimal Controls in the Absence of Convexity Condition", Jour. of Math. Ana. and App. (to appear) 1963.